## **ACT-FAST**

ACT-FAST is a large vessel occlusion screening tool utilized for anterior circulation strokes. It is a two step process that provides a positive or negative result by assessing for one-sided weakness, language deficit, gaze preference or hemi-neglect.

Step three guides clinical decision making in the eligibility for Endovascular Treatment (EVT). Registered Nurses, Registered Practical Nurses, Nurse Practitioners, Physician Assistants and Physicians can preform the ACT-FAST screening

### **ACT-FAST Assessment**

### **Step One**

#### Proceed if POSTIVE

If negative STOP

Notify ED Physician and continue with usual care



### ARM Unilateral Arm Weakness

Position both arms with elbows straight (45 degrees if supine, 90 degrees if sitting)

#### **Positive Screen:**

- · One arm falls completely within 10 seconds
- · For patients who are uncooperative or cannot follow commands: if you witness minimal or no movement in one arm and normal movement in the other

### **Step Two**



Proceed if POSTIVE

If negative STOP

Notify ED Physician and continue with usual care

### Unilateral **RIGHT** arm weakness?



# CHAT Severe Language Deficit

Ask patient to repeat a phrase or perform simple tasks

### **Positive Screen**:

- · Patient is mute, speaking gibberish and/or incomprehensible
- · Patient is unable to follow simple commands

### Unilateral **LEFT** arm weakness?

# TAP Gaze and Shoulder Tap Test



Stand on patients weak side while assessing

#### **Positive Screen:**

- Patient has consistent gaze away from weak side
- Patient does not quickly turn head/eyes to you when shoulder tapped twice and first name called

### **Step Three**



If POSTIVE, potential LVO

### **Eligibility for EVT**

**Positive screen:** (if all criteria met)

- Deficits are NOT pre-existing (mild deficits now worse are acceptable as true deficits)
- Living at home independently with only minor assistance must be independent with hygiene, personal care, walking (walking aids not included)
- Does NOT have stroke mimics: seizure preceding symptoms, hypoglycemia (glucose less than 2.8 mmol/L), active malignancy with brain lesions



Acronyms: ACT-FAST Ambulance Clinical Triage for Acute Stroke Treatment Source: strokebestpractices.ca v0.1 Sept 2022

